


Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



(This is a sample cover image for this issue. The actual cover is not yet available at this time.)

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the author's institution and sharing with colleagues.


Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Toxicon

journal homepage: <http://www.elsevier.com/locate/toxicon>

Short communication

19-Hydroxy-bufalin, a major bufadienolide isolated from the parotoid gland secretions of the Panamanian endemic toad *Rhinella centralis* (Bufonidae), inhibits the growth of *Trypanosoma cruzi*

Candelario Rodríguez^{a,b,c}, Roberto Ibáñez^{d,e}, Michelle Ng^f, Carmenza Spadafora^f, Armando A. Durant-Archibold^{a,c}, Marcelino Gutiérrez^{a,*}

^a Centro de Biodiversidad y Descubrimiento de Drogas, Instituto de Investigaciones Científicas y Servicios de Alta Tecnología (INDICASAT AIP), Panama

^b Department of Biotechnology, Acharya Nagarjuna University, India

^c Departamento de Bioquímica, Facultad de Ciencias Naturales, Exactas y Tecnología, Universidad de Panamá, Panama

^d Smithsonian Tropical Research Institute (STRI), Panama

^e Departamento de Zoología, Facultad de Ciencias Naturales, Exactas y Tecnología, Universidad de Panamá, Panama

^f Centro de Biología Celular y Molecular de Enfermedades, INDICASAT AIP, Panama



ARTICLE INFO

Keywords:

Rhinella centralis

Toad

Bufonidae

Parotoid gland secretion

Bufadienolide

Antitrypanosomal activity

Trypanosoma cruzi

ABSTRACT

American trypanosomiasis is a parasitic neglected disease, responsible for the death of approximately 10,000 people every year. Amphibians are recognized for producing in their cutaneous glands substances with pharmacological potential against a variety of pathologies. Here we investigated the antiprotozoal activity against *Trypanosoma cruzi* of bufadienolides isolated from the parotoid glands secretions of the toad *Rhinella centralis* from Panama. NMR and mass spectrometry analysis led to the identification of the active compound 19-hydroxy-bufalin, for which its antitrypanosomal activity and occurrence in the genus *Rhinella* are reported for the first time. This compound showed low cytotoxicity and significant selectivity which confers to it a potential role for the treatment of Chagas disease.

American trypanosomiasis, also known as Chagas disease, is a zoonotic disease caused by the protozoan parasite *Trypanosoma cruzi*. Each year more than 10,000 people worldwide die of Chagas disease-related causes, and 8 million people are estimated to be infected by *T. cruzi* in Latin America (WHO, 2019). Currently, benznidazole and nifurtimox are the drugs available for the treatment of Chagas disease; however, these medications lead to the development of severe secondary effects and are effective only during the first weeks after infection (Castro et al., 2006).

Amphibians of the genus *Rhinella* contains in their skin secretions metabolites of the bufadienolides family, which are polyhydroxy steroids containing a 2-pyrone ring at position 17 of the steroidal core. These compounds are of interest due to their high chemical diversity and promising pharmacological activities against bacteria, fungi and trypanosomatid parasites (Barnhart et al., 2017; Cunha Filho et al., 2005; Tempone et al., 2008). The endemic Panamanian toad *Rhinella centralis* (Narvaes and Rodrigues, 2009), formerly known as *Bufo granulosus*,

inhabits the lowlands of the Pacific slope of Panamá from Chiriquí to Darién provinces (Frost, 2019; IUCN, 2019). This relatively small species of toad is nocturnal and terrestrial, and breeds throughout the wet season particularly on rainy nights (Roberto Ibáñez et al., 1999). In this communication, we are reporting for the first time the occurrence in the genus *Rhinella* and the antitrypanosomal activity of 19-hydroxy-bufalin, a bufadienolide isolated from the parotoid gland secretions of the toad *R. centralis*.

Adult specimens of *R. centralis* were collected in Pacora, Panamá Province (Fig. 1-A) (collection permit SE/A-32-15). Parotoid glands secretion was obtained by electrical stimulation, after applying 5 V at 60 Hz for 30 s on the toads' dorsal region, using a homemade transcutaneous amphibian stimulator (Grant and Land, 2002). The secretions were placed in vials containing MeOH and stored at -20°C until analysis. The methanolic soluble extract was loaded in Supelclean LC-18 solid phase extraction (SPE) cartridges (Supelco, Bellefonte, PA, USA), and fractionated using vacuum employing a three-step gradient

* Corresponding author. Centro de Biodiversidad y Descubrimiento de Drogas, Instituto de Investigaciones Científicas y Servicios de Alta Tecnología (INDICASAT AIP), Panamá, Apartado, 0843-01103, Panama.

E-mail address: mgutierrez@indicat.org.pa (M. Gutiérrez).

<https://doi.org/10.1016/j.toxicon.2020.02.009>

Received 5 December 2019; Received in revised form 3 February 2020; Accepted 11 February 2020

Available online 13 February 2020

0041-0101/© 2020 Elsevier Ltd. All rights reserved.

composed of mixtures of 40, 80 and 100% of methanol in water at 10 mL min^{-1} . Methanol was removed by rotary evaporation under reduced pressure and the remaining water lyophilized obtaining three fractions F1–F3. The three fractions were evaluated against trypomastigotes of *T. cruzi*. Active fraction F2 was diluted in MeOH at 30 mg mL^{-1} and subjected to reversed-phase high performance liquid chromatography (RP-HPLC) employing a semi-preparative Polar C18 column ($250 \times 10 \text{ mm}$, $4 \mu\text{m}$, 80 \AA) (Phenomenex®, Torrance, CA, USA). Injections of $100 \mu\text{L}$ were carried out and elution was set at a flow rate of 3 mL min^{-1} . The solvent system consisted of water with 0.1% of trifluoroacetic acid (TFA) (Solvent A), and a 1/1 mixture of MeOH/ CH_3CN acidified with TFA at 0.1% (Solvent B). Elution was carried out using a gradient from 20% to 100% of solvent B in solvent A in 60 min.

Structural elucidation was carried out by spectroscopic and spectrometric means. Nuclear magnetic resonance (NMR) experiments were carried out at $25 \text{ }^\circ\text{C}$ on an Eclipse+ 400 Fourier transform spectrometer (JEOL, Peabody, MA, USA). For NMR analysis, the samples were dissolved in deuterated methanol, and the spectra referenced based on the non-deuterated methanol residual signal (3.31 for δ_{H} and 49.0 for δ_{C}). High resolution mass spectrometry analysis was carried out on an UltrafleXtreme MALDI-TOF/TOF (Bruker Daltonics, Billerica, MA, USA). For MALDI-TOF-TOF analysis, $2 \mu\text{L}$ of the sample (19-hydroxy-bufalin) dissolved in methanol, were transferred to a MTP plate (Bruker Daltonics, Billerica, MA, USA), and mixed with $1 \mu\text{L}$ of the matrix (α -cyano-4-hydroxy-cinammic acid, Sigma Aldrich). All experiments were carried out in reflectron positive mode. Peptide calibration standard II (Bruker, Daltonics, MA) was used as external calibrant.

Antitrypanosomal bioassay was carried out on recombinant Tulahuén clone C4 of *T. cruzi* trypomastigotes (American Type Culture Collection, ATCC), Manassas, VA, USA), which expresses the

β -galactosidase enzyme that serves as a reporter of viability (Buckner et al., 1996). Transfected parasites were maintained at $37 \text{ }^\circ\text{C}$ under a 5% CO_2 atmosphere in RPMI-1640 culture medium with L-glutamine, 4-(2-Hydroxyethyl)-piperazine-1-ethanesulfonic acid buffer, NaHCO_3 , 10% FBS and 1% penicillin/streptomycin as supplements.

Vero cells (ATCC) were grown for 24 h, and one day prior to the experiment the cells were infected with trypomastigotes. Samples were dissolved in DMSO and added to the medium at 10.0, 2.0, 0.4 and $0.1 \mu\text{g/mL}$ and incubated for five days. To determine the antitrypanosomal activity, chlorophenol-red- β -D-galactopyranoside (Roche Applied Science) was added to each well and allowed to react with the β -galactosidase of the remnant living parasites for 4.5 h. Benznidazole was used as positive control ($\text{IC}_{50} = 0.75 \mu\text{g/mL}$). Absorbance was measured at 570 nm using a color plate reader (Sinergy HT, BioTek Instruments Inc, Winooski, VT, USA).

For cytotoxicity assays, Vero cells were cultivated in 96-well plates at $37 \text{ }^\circ\text{C}$ under a 5% CO_2 atmosphere, using RPMI-1640 medium (Sigma-Aldrich, USA) sterilized with 1% penicillin/streptomycin and supplemented with 10% FBS (fetal bovine serum; Gibco, Invitrogen, Carlsbad, CA, USA). The cells were allowed to adhere for 24 h prior to incubation for five days with the samples. During the treatment, DMSO was used as negative control, and samples were assayed at the same concentrations used in the anti-trypanosomal assay. After incubation, MTT (3-(4, 5-dimethyl-thiazol-2-yl)-2, 5-di-phenyl-tetra-zolium bromide) was added to each well and absorbance was determined 4 h later at 570 nm, using a color plate reader. Cytotoxicity was evaluated colorimetrically by calculating the ability of the remnant living Vero cells to reduce the pale yellow MTT into the black purple formazan product, as described earlier (Mosmann, 1983). All bioassays were performed in duplicate and the IC_{50} values were determined employing the Data Analysis complement




Fig. 1. (A) Distribution range, collection site and adult of *Rhinella centralis*. Its distribution was partly based on IUCN. (2019). (B) RP-HPLC Chromatogram profile of the fraction F2 obtained by VLC (80% MeOH in H_2O) from parotoid glands secretion of *R. centralis*. Black arrow indicates 19-hydroxy-bufalin (1). (C) Key COSY and HMBC correlations of 19-hydroxy-bufalin (1). (D) Effect of 19-hydroxy-bufalin (1) against trypomastigotes of *T. cruzi*.

Wizard of Excel (2000) (Microsoft, Seattle, WA, USA).

The skin secretion of *R. centralis* (1093 mg) subjected to SPE fractionation resulted in three fractions F1–F3. Antitrypanosomal activity was found in fraction F2 (320 mg) which was subjected to semi-preparative RP-HPLC purification leading to the isolation of compound **1** (17.6 mg), eluted at 27.7 min, active against *T. cruzi* trypomastigotes (Fig. 1-B). The structural elucidation of compound **1** was carried out by spectroscopic analyses and comparison with data from the literature (Verpoorte and Svendsen, 1979).

The molecular formula of compound **1** was determined as C₂₄H₃₄O₅ based on HR-MALDI-TOF-TOF MS data which showed a sodiated adduct of the molecular ion at *m/z* 425.2295 [M + Na]⁺ (calcd for C₂₄H₃₄O₅Na, 425.2298). Eight degrees of unsaturation were calculated from the molecular formula: three accounted for a carbonyl and two double bonds, hence compound **1** possess five rings.

In general, analysis of the ¹H and ¹³C NMR data (Table 1) revealed the typical spectroscopic features of a bufadienolide including resonances for the 2-pyrone ring at δ_C 125.1 (C-20), 150.5 (C-21), 149.5 (C-22), 115.56 (C-23) and 164.9 (C-24), (Verpoorte et al., 1980). ¹H NMR showed signals for two diastereotopic protons mutually coupled, consistent with an oxomethylene at δ_H 3.83 (d, *J* = 11.2 Hz) and 3.41 (d, *J* = 11.2 Hz), attached to carbon C-19 (δ_C 66.17) using HSQC data. The steroidal nucleus was assembled using COSY and *J*^{2,3} HMBC correlations (Fig. 1-C) being consistent with the structure of 19-hydroxy-bufalin (**1**).

19-hydroxy-bufalin (**1**) has been isolated from toads of the genera *Bufo*, *Bufo*tes and *Duttaphrynus*, although not known in the genus *Rhinella* (Gao et al., 2010; Gella et al., 1995; Verpoorte et al., 1980). In our study, 19-hydroxy-bufalin (**1**) led to a growth inhibition in a dose-dependent manner with an IC₅₀ of 7.81 μg/mL (Fig. 1-D) when assayed against *T. cruzi* trypomastigotes. Moreover, the cytotoxicity evaluation on epithelial kidney Vero cells exhibited a low inhibitory concentration with an IC₅₀ of 71.58 μg/mL, resulting in a significant selectivity index (SI = IC₅₀ Vero cells/IC₅₀ *T. cruzi*) of 9.16. The selectivity index is a good reference of the pharmacological activity and safety of a drug. For instance, the higher the SI, the more effective and safe would be a compound during *in vivo* antiprotozoal treatment (Nogueira et al., 2009).

Only one bufadienolide, hellebrigenin, isolated from the parotoid macroglands of *Rhinella jimi*, has shown activity against *T. cruzi*, with a reported IC₅₀ of 91.75 μg/mL (Tempone et al., 2008). To the best of our knowledge, in this study we report for the first time the anti-trypomastigote activity of 19-hydroxy-bufalin and its occurrence in the genus *Rhinella*. The therapeutic window of 19-hydroxy-bufalin against American trypanosomiasis points out the need for research on the mechanism by which this natural product acts. This communication highlights the importance of amphibians as a source for the discovery of natural products with potential for the treatment of neglected human diseases.

Ethical statement

Marcelino Gutiérrez and the coauthors state that this work submitted here has not been submitted for publication elsewhere. All work was conducted to the ethical standards outlined in the Publishing Ethics Resource Kit associated found on the journal webpage.

Declaration of competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

CRediT authorship contribution statement

Candelario Rodriguez: Data curation, Formal analysis, Investigation, Methodology, Writing - original draft, Writing - review & editing.

Table 1

¹H and ¹³C NMR data (400 MHz) of 19-hydroxy-bufalin (**1**) in CD₃OD, chemical shifts (δ) in ppm.

No.	δ _C , type ^a	δ _H , mult. ^{b,c,d} (<i>J</i> in Hz)
1	24.3, CH ₂	1.88, m 1.39, m
2	33.1, CH ₂	2.13, m 1.72, m
3	67.6, CH	4.04, m
4	28.4, CH ₂	1.58, m n.d.
5	29.8, CH	2.21, m
6	27.6, CH ₂	1.88, m 1.22, m
7	22.8, CH ₂	1.49, m 1.22, m
8	43.0, CH	1.68, m
9	36.6, CH	1.80, m
10	40.6, C	
11	22.3, CH ₂	1.80, m 1.35, m
12	42.4, CH ₂	1.49, m 1.39, m
13	49.9, C	
14	86.4, C	
15	34.2, CH ₂	1.85, m 1.39, m
16	29.9, CH ₂	2.13, m 1.72, m
17	52.4, CH	2.54, m
18	17.4, CH ₃	0.70, s
19	66.1, CH ₂	3.83, d (11.24) 3.41, d (11.24)
20	125.1, C	
21	150.5, CH	7.42, m
22	149.5, CH	7.98, dd (2.44, 9.28)
23	115.5, CH	6.27, d (9.76)
24	164.9, C	

^a Type of carbon was determined by DEPT spectra (90° and 135°).

^b Assignments were made by analysis of COSY, HSQC and HMBC spectra.

^c Multiplets are identified as s (singlet), d (doublet), dd (doublet of doublets) or m (when multiplicity is not clear).

^d Overlapped signals are reported without multiplicity. n.d. = not detected.

Roberto Ibáñez: Data curation, Formal analysis, Investigation, Methodology, Resources, Supervision, Writing - original draft, Writing - review & editing. **Michelle Ng:** Data curation, Formal analysis, Investigation, Methodology. **Carmenza Spadafora:** Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Resources, Supervision, Writing - original draft, Writing - review & editing. **Armando A. Durant-Archibold:** Conceptualization, Data curation, Formal analysis, Investigation, Resources, Supervision, Methodology, Writing - original draft, Writing - review & editing. **Marcelino Gutiérrez:** Conceptualization, Formal analysis, Funding acquisition, Investigation, Project administration, Resources, Supervision, Validation, Methodology, Data curation, Writing - original draft, Writing - review & editing.

Acknowledgments

CR was supported by a scholarship from the Instituto para la Formación y Aprovechamiento de Recursos Humanos (IFARHU), Panama and The Secretaría Nacional de Ciencia, Tecnología e Innovación of Panama (SENACYT), Panama. We gratefully acknowledge the Government of Panama (MiAmbiente) for granting permission to collect the frogs used in this study. The authors acknowledge SENACYT for financial support through the INDICASAT-BID Program (grant numbers IND-JAL-05 and 02-12-H). CR, RI, AAD-A, CS and MG were supported by the Sistema Nacional de Investigación (SNI), Panama. RI was supported by the Panama Amphibian Rescue and Conservation Project and Minera Panamá. CS was supported by grant FID17-095 from SENACYT. We

thank Arellys Fuentes for her assistance obtaining the HRMS data.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.toxicol.2020.02.009>.

References

- Barnhart, K., Forman, M.E., Umile, T.P., Kueneman, J., McKenzie, V., Salinas, I., Minbiole, K.P.C., Woodhams, D.C., 2017. Identification of bufadienolides from the boreal toad, *Anaxyrus boreas*, active against a fungal pathogen. *Microb. Ecol.* 74, 990–1000. <https://doi.org/10.1007/s00248-017-0997-8>.
- Buckner, F.S., Verlinde, C.L.M.J., Flamme, A.C.L.A., 1996. Efficient technique for screening drugs for activity against *Trypanosoma cruzi* using parasites expressing beta-galactosidase, 40, 2592–2597. <https://doi.org/10.1128/AAC.40.11.2592>.
- Castro, J.A., De Mecca, M.M., Bartel, L.C., 2006. Toxic side effects of drugs used to treat Chagas' disease (American trypanosomiasis). *Hum. Exp. Toxicol.* 25, 471–479. <https://doi.org/10.1191/0960327106het653oa>.
- Cunha Filho, G.A., Schwartz, C.A., Resck, I.S., Murta, M.M., Lemos, S.S., Castro, M.S., Kyaw, C., Pires, O.R., Leite, J.R.S., Bloch, C., Schwartz, E.F., 2005. Antimicrobial activity of the bufadienolides marinobufagin and telocinobufagin isolated as major components from skin secretion of the toad *Bufo rubescens*. *Toxicol* 45, 777–782. <https://doi.org/10.1016/j.toxicol.2005.01.017>.
- Frost, D.R., 2019. Amphibian Species of the World: an Online Reference. American Museum of Natural History, New York, USA (13 November 2019). Electronic Database accessible at, Version 6.0. <http://research.amnh.org/herpetology/amphibia/index.html>.
- Gao, H., Zehl, M., Leitner, A., Wu, X., Wang, Z., Kopp, B., 2010. Comparison of toad venoms from different *Bufo* species by HPLC and LC-DAD-MS/MS. *J. Ethnopharmacol.* 131, 368–376. <https://doi.org/10.1016/j.jep.2010.07.017>.
- Gella, I.M., Shabanov, D.A., Leont'ev, D.A., Levin, M.G., Shishkin, O.V., Baumer, V.N., Lakin, E.E., 1995. Study of the bufadienolides of the skin secretion of green toads (*Bufo viridis* laur, 1758). *Pharm. Chem. J.* 29, 491–494. <https://doi.org/10.1007/BF02220008>.
- Grant, J.B., Land, B., 2002. Transcutaneous Amphibian Stimulator (TAS): a device for the collection of amphibian skin secretions. *Herpetol. Rev.* 33, 38–41.
- IUCN, 2019. The IUCN red list of threatened species. Available online. <https://www.iucnredlist.org/search?query=rhinella%20centralis&searchType=species>. (Accessed 13 November 2019). accessed on.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* 65, 55–63. [https://doi.org/10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4).
- Narvaes, P., Rodrigues, M.T., 2009. Taxonomic revision of *Rhinella granulosa* species group (Amphibia, Anura, Bufonidae), with a description of a new species. *Arq. Zool. (Sao Paulo)* 40, 1. <https://doi.org/10.11606/issn.2176-7793.v40i1p1-73>.
- Nogueira, R.C., Oliveira-Costa, J.F., de Sá, M.S., dos Santos, R.R., Soares, M.B.P., 2009. Early toxicity screening and selection of lead compounds for parasitic diseases. *Curr. Drug Targets* 10, 291–298.
- Roberto Ibáñez, D., Stanley Rand, A., Jaramillo, César A., 1999. A. Los anfibios del Monumento Natural Barro Colorado, Parque Nacional Soberanía y sus áreas adyacentes, first ed.
- Tempone, A.G., Pimenta, D.C., Lebrun, I., Sartorelli, P., Taniwaki, N.N., de Andrade, H. F., Antoniazzi, M.M., Jared, C., 2008. Antileishmanial and antitrypanosomal activity of bufadienolides isolated from the toad *Rhinella jimi* parotoid macrogland secretion. *Toxicol* 52, 13–21. <https://doi.org/10.1016/j.toxicol.2008.05.008>.
- Verpoorte, R., Svendsen, A.B., 1979. Chemical constituents of Vietnamese toad venom collected from *Bufo melanostictus* schneider. *J. Ethnopharmacol.* 1, 197–202. [https://doi.org/10.1016/0378-8741\(79\)90007-2](https://doi.org/10.1016/0378-8741(79)90007-2).
- Verpoorte, R., Phax-quôc-Kixh, Svendsen, A.B., 1980. Chemical constituents of Vietnamese toad venom, collected from *Bufo melanostictus* schneider. Part II. The bufadienolides. *J. Nat. Prod.* 43, 347–352. <https://doi.org/10.1021/np50009a005>.
- WHO, World Health Organization, 2019. Available online. <https://www.who.int/chagas/en/>. (Accessed 13 November 2019). accessed on.